ELSEVIER

Biomembranes

Vol. 1416, Nos. 1–2; 12 January 1999

BIOCHIMICA ET BIOPHYSICA ACTA

BBA

Contents

BBA is cited in: Current Contents/Life Sciences - Biological Abstracts - Chemical Abstracts - Index Chemicus - Index Medicus/MEDLINE - Cambridge Scientific Abstracts (CSA) - EMBASE/Excerpta Medica - Reference Update - Elsevier BIOBASE/Current Awareness in Biological Sciences

Information for Contributors *vii*

Regular papers

Factors influencing uptake and retention of amino-containing drugs in large unilamellar vesicles exhibiting transmembrane pH gradients
E. Maurer-Spurej, K.F. Wong, N. Maurer, D.B. Fenske and P.R. Cullis (Canada) 1

Palmitoylation of lung surfactant protein SP-C alters surface thermodynamics, but not protein secondary structure or orientation in 1,2-dipalmitoylphosphatidylcholine Langmuir films
C.R. Flach, A. Gericke, K.M.W. Keough and R. Mendelsohn (Canada, Germany, USA) 11

Surface charge response of the phosphatidylcholine head group in bilayered micelles from phosphorus and deuterium nuclear magnetic resonance
K.J. Crowell and P.M. Macdonald (Canada) 21

Different kinds of polypeptides and polypeptide-coated nanoparticles are accepted by the selective transcytosis shown in the rabbit nasal mucosa
D. Cremaschi, C. Porta and R. Ghirardelli (Italy) 31

Identification of particular epithelial areas and cells that transport polypeptide-coated nanoparticles in the nasal respiratory mucosa of the rabbit
R. Ghirardelli, F. Bonasoro, C. Porta and D. Cremaschi (Italy) 39

NMR structure of C-terminally tagged gramicidin channels
F. Separovic, S. Barker, M. Delahunty and R. Smith (Australia) 48

Enzymatic reactions in liposomes using the detergent-induced liposome loading method
T. Oberholzer, E. Meyer, I. Amato, A. Lustig and P.-A. Monnard (Switzerland) 57

Two-component blocking kinetics of open NMDA channels by organic cations
A.I. Sobolevsky (Russia) 69

Functional reconstitution of bacterially expressed human potassium channels in proteoliposomes: membrane potential measurements with JC-1 to assay ion channel activity
B. Chanda and M.K. Mathew (India) 92

Cleavage of annexin I in human neutrophils is mediated by a membrane-localized metalloprotease
C. Movitz, C. Sjölin and C. Dahlgren (Sweden) 101

Nucleobase transport in opossum kidney epithelial cells and *Xenopus laevis* oocytes: the characterisation, structure–activity relationship of uracil analogues and oocyte expression studies of sodium-dependent and -independent hypoxanthine uptake
M. Shayeghi, R. Akerman and S.M. Jarvis (UK) 109

Calorimetric and spectroscopic studies of the effects of cholesterol on the thermotropic phase behavior and organization of a homologous series of linear saturated phosphatidylethanolamine bilayers
T.P.W. McMullen, R.N.A.H. Lewis and R.N. McElhaney (Canada) 119

Determination by photoreduction of flip-flop kinetics of spin-labeled stearic acids across phospholipid bilayers
J.-M.P. Yuann and R.(P.D.II) Morse (USA) 135

Glucose transporter 3 (GLUT3) protein is present in human myocardium
M. Grover-McKay, S.A. Walsh and S.A. Thompson (USA) 145

Identification of 130 kDa cell surface LDL-binding protein from smooth muscle cells as a partially processed T-cadherin precursor
D.V. Stambolsky, Y.S. Kuzmenko, M.P. Philippova, V.N. Bochkov, Z.D. Bespalova, A.A. Azmuko, N.M. Kashirina, T.N. Vlasik, V.A. Tkachuk and T.J. Resink (Russia, Switzerland) 155

The amphipathic helix concept: length effects on ideally amphipathic $L_iK_j(i=2j)$ peptides to acquire optimal hemolytic activity
S. Castano, I. Cornut, K. Büttner, J.L. Dasseux and J. Dufourcq (France, Germany) 161

Structure, orientation and affinity for interfaces and lipids of ideally amphipathic lytic $L_iK_j(i=2j)$ peptides
S. Castano, B. Desbat, M. Laguerre and J. Dufourcq (France) 176

Activation of Na^+,K^+,Cl^- cotransport in squid giant axon by extracellular ions: evidence for ordered binding
A.A. Altamirano, G.E. Breitwieser and J.M. Russell (USA) 195